


Using the Shell Method for Finding Rotational Volumes

W-up: AP Multiple Choice #86(graphing calculator allowed)


The Shell Method: Used when it is convenient to represent the radius *in terms of* x but rotation is about a vertical axis(like the y -axis) or vice-versa. Also NEEDED when an equation cannot be solved **for a specific variable**.

EX) Find the volume of a solid of revolution formed by revolving the 1st quadrant region bounded by $f(x) = x - x^3$ and $y = 0$ about the y -axis (to be solved later).

The shell method uses the lateral area of cylinders to find volumes



$$\text{Lateral Area} = 2\pi rh$$



Shell Method for **Vertical** Axis of Revolution

$$V = 2\pi \int_a^b p(x) \cdot h(x) dx$$

Where $p(x)$ represents the algebraic expression for the radius of each cylinder(distance between the *axis of revolution* and any “ x ”) on $[a, b]$ AND $h(x)$ represents the algebraic expression for the height of each cylinder (distance between boundaries of the bounded region) on $[a, b]$.

Shell Method for **Horizontal** Axis of Revolution

$$V = 2\pi \int_c^d p(y) \cdot h(y) dy$$

Where $p(y)$ represents the algebraic expression for the radius of each cylinder (distance between the *axis of revolution* and any “y”) on $[c, d]$ AND $h(y)$ represents the algebraic expression for the height of each cylinder (distance between boundaries of the bounded region) on $[c, d]$.

Remember: When finding vertical distance (which is always positive) on the coordinate plane use the “top value – bottom value” and when finding horizontal distance use the “right value – left value.”

EX) Find the volume of a solid of revolution formed by revolving the 1st quadrant region bounded by $f(x) = x - x^3$ and $y = 0$ about the y-axis

EX) Find the volume of a solid of revolution formed by revolving the 1st quadrant region bounded by $f(x) = 2x^2 + 3$, $x = 0$ and $y = 7$ about the y-axis